

**Association between psychotic experiences and subsequent suicidal thoughts and behaviors: a cross-national analysis from the World Health Organization World Mental Health Surveys**

Bromet EJ, Nock MK, Saha S, Lim CCW, Aguilar-Gaxiola S, Al-Hamzawi A, Alonso J, Borges GLG, Bruffaerts R, Degenhardt L, de Girolamo G, de Jonge P, Florescu S, Gureje O, Haro JM, He Y, Hu C, Karam EG, Kovess-Masféty V, Lee S, Lépine JP, Mneimneh Z, Navarro-Mateu F, Ojagbemi A, Posada-Villa J, Sampson NA, Scott KM, Stagnaro JC, Viana MC, Xavier M, Kessler RC, McGrath JJ.

JAMA Psychiatry

2017; 74(11):1136-1144

**ARTICLE IDENTIFIERS**

DOI: [10.1001/jamapsychiatry.2017.2647](https://doi.org/10.1001/jamapsychiatry.2017.2647)

PMID: 28854302

PMCID: not available

**JOURNAL IDENTIFIERS**

LCCN: 2012200352

pISSN: 2168-622X

eISSN: 2168-6238

OCLC ID: 798278977

CONS ID: not available

US National Library of Medicine ID: 101589550

This article was identified from a query of the SafetyLit database.