

## **Risk of depression, suicide and psychosis with hydroxychloroquine treatment for rheumatoid arthritis: a multinational network cohort study**

Lane JCE, Weaver J, Kostka K, Duarte-Salles T, Abrahao MTF, Alghoul H, Alser O, Alshammari TM, Areia C, Biedermann P, Banda JM, Burn E, Casajust P, Fister K, Hardin J, Hester L, Hripcsak G, Kaas-Hansen BS, Khosla S, Kolovos S, Lynch KE, Makadia R, Mehta PP, Morales DR, Morgan-Stewart H, Mosseveld M, Newby D, Nyberg F, Ostropolets A, Woong Park R, Prats-Urbe A, Rao GA, Reich C, Rijnbeek P, Sena AG, Shoaibi A, Spotnitz M, Vignesh S, Suchard MA, Vizcaya D, Wen H, de Wilde M, Xie J, You SC, Zhang L, Lovestone S, Ryan P, Prieto-Alhambra D.

Rheumatology (Oxford)

2020; ePub(ePub):ePub

### **ARTICLE IDENTIFIERS**

DOI: [10.1093/rheumatology/keaa771](https://doi.org/10.1093/rheumatology/keaa771)

PMID: unavailable

PMCID: not available

### **JOURNAL IDENTIFIERS**

LCCN: sn 99047125

pISSN: 1462-0324

eISSN: 1462-0332

OCLC ID: 40922029

CONS ID: not available

US National Library of Medicine ID: 100883501

This article was identified from a query of the SafetyLit database.